

SOLUTIONS & ANSWERS FOR JEE MAINS-2016 VERSION – E

[PHYSICS, CHEMISTRY & MATHEMATICS]

PART – A – PHYSICS

1. A student measures the time period of 100 oscillations ----

Ans: 92 ± 2 s

Sol:
$$\bar{x} = \frac{\sum x_i}{N} = 92$$

$$\Delta \bar{x} = \frac{\sum \Delta x}{N} = 1.5 \text{ s}$$

The least count is 1 s, $\Delta x = 2$ s
Reported mean time = 92 ± 2 s


2. A particle of mass m is moving along the side of a square of side ----

Ans:
$$\vec{L} = mv \left[\frac{R}{\sqrt{2}} - a \right] \hat{k}$$
 when the particle is moving from, C to D.

AND

$$\vec{L} = \frac{mv}{\sqrt{2}} R \hat{k}$$
 when the particle is moving from, D to A.

Sol:



$$\vec{L}_{AB} = mv \frac{R}{\sqrt{2}} (-\hat{k})$$
 [at A]

$$\vec{L}_{CD} = mv \left(\frac{R}{\sqrt{2}} + a \right) (+\hat{k})$$
 [at C]

$$\vec{L}_{BC} = mv \left(\frac{R}{\sqrt{2}} + a \right) (+\hat{k})$$
 [at C]

$$\vec{L}_{DA} = mv \frac{R}{\sqrt{2}} (-\hat{k})$$
 [at A]

Option (2 and 4) can be the answer.

3. A point particle of mass m , moves along the uniformly ----

Ans: 0.29 and 3.5 m

Sol: $\mu mg \cos \theta \times PQ = \mu mg QR$ (data)

$$\frac{x}{PQ} \times PQ = QR \Rightarrow x = QR$$

$$\tan 30^\circ = \frac{2}{x} = \frac{1}{\sqrt{3}} \Rightarrow QR = x = 2\sqrt{3}$$

= 3.46 m

= 3.5 m

Work energy theorem

$-\mu mgx - \mu mg(QR) = 0 - mgh$

$2\mu x = h$

$$\mu = \frac{h}{2x} = \frac{2}{2 \times 2\sqrt{3}} = \frac{1}{2\sqrt{3}} = \frac{\sqrt{3}}{6}$$

= $\frac{1.732}{6} = 0.3$

4. A person trying to lose weight by burning fat lifts a mass of 10 kg ----

Ans: 12.89×10^{-3}

Sol: $m \times 3.8 \times 10^7 \times 20\%$
= $10 \times 9.8 \times 1 \times 1000$


$$m = \frac{9.8 \times 10^4}{3.8 \times 0.2 \times 10^7}$$

= 12.89×10^{-3} kg

5. A roller is made by joining together two cones at their vertices O. It is kept ----

Ans: turn left

Sol:



At point of contact C_1 $v = \omega R$ (pure roll)

At point of contact C_2 $v > \omega r \Rightarrow$

forward slipping, \therefore friction will act backwards towards C_1 .

Roller will turn anti-clockwise since torque of friction about CM is anti-clockwise.

6. A satellite is revolving in a circular orbit at a height 'h' from the earth's ----

Ans: $\sqrt{gR}(\sqrt{2} - 1)$

Sol: $v_{es} = \sqrt{\frac{2GM}{R}} = \sqrt{2gR}$
 $v_0 = \sqrt{\frac{GM}{R}} = \sqrt{gR}$
 Increase = $\sqrt{2gR} - \sqrt{gR} = \sqrt{gR}[\sqrt{2} - 1]$

7. A pendulum clock loses 12 s a day if the temperature is 40 °C and gains 4 s ----

Ans: 25 °C; $\alpha = 1.85 \times 10^{-5} / ^\circ\text{C}$

Sol: $12 = \frac{1}{2} \alpha(40 - T_0)$

$4 = \frac{1}{2} \alpha(T_0 - 20)$

If temperature increases, clock loses time and temperature decreases, clock gains time.

Solving $T_0 = 25 ^\circ\text{C}$, $\alpha = 1.85 \times 10^{-5} / ^\circ\text{C}$

8. An ideal gas undergoes a quasi static, reversible process in which its ----

Ans: $n = \frac{C - C_p}{C - C_v}$

Sol: $pV^n = \text{constant}$, $\mu = \text{no. of moles}$

work = $\int pdV = \frac{\mu R \Delta T}{-n+1}$

According to I law of thermodynamics

$\mu C \Delta T = \mu C_v \Delta T + \frac{\mu R \Delta T}{-n+1}$

$C - C_v + \frac{C_p - C_v}{-n+1}$

$-n + 1 = \frac{C_p - C_v}{C - C_v}$


$n = 1 - \frac{C_p - C_v}{C - C_v} = \frac{C - C_v - C_p + C_v}{C - C_v}$

$= \frac{C - C_p}{C - C_v}$

9. 'n' moles of an ideal gas undergoes a process A → B as ----

Ans: $\frac{9 p_0 V_0}{4 nR}$

Sol: Draw the isotherm through A & B ($2p_0V_0 = nRT$)



Maximum temperature will be at mid point of AB; that is an isotherm tangent to the straight line AB at its mid point.

$p = 1.5p_0$ $V = 1.5V_0$

$pV = nRT \Rightarrow T = \frac{1.5p_0 \times 1.5V_0}{nR}$

$= \frac{9 p_0 V_0}{4 nR}$

10. A particle performs simple harmonic motion with amplitude A. Its speed is ----

Ans: $\frac{7}{3} A$

Sol: Potential energy $U = \frac{1}{2} K \left(\frac{2}{3} A \right)^2$

Kinetic energy = $\frac{1}{2} K \left[A^2 - \left(\frac{2}{3} A \right)^2 \right]$

New K.E = 9K (K.E $\propto v^2$)

New total energy = $9K + U = \frac{1}{2} K A_1^2$


$\frac{9}{2} K \left[A^2 - \frac{4}{9} A^2 \right] + \frac{1}{2} K \frac{4}{9} A^2$

$= \frac{1}{2} K A_1^2 \Rightarrow A_1 = \frac{7}{3} A$

11. A uniform string of length 20 m is suspended from a rigid support. A short ----

Ans: $2\sqrt{2}$ s

Sol:



$\mu = \text{mass per unit length}$

$v = \sqrt{\frac{T}{\mu}} = \sqrt{\frac{\mu x g}{\mu}}$

$$\Rightarrow v = \sqrt{xg} = \frac{dx}{dt}$$

$$\int \sqrt{g} dt = \int_0^{20} x^{-\frac{1}{2}} dx$$

$$\sqrt{g} \times t = \frac{x^{-\frac{1}{2}+1}}{-\frac{1}{2}} = 2\sqrt{x} \Big|_0^{20}$$

$$\sqrt{10} \times t = 2 \times \sqrt{20}$$

$$\Rightarrow t = 2\sqrt{2} \text{ s}$$

12. The region between two concentric spheres of radii 'a' and 'b', respectively ----

Ans: $\frac{Q}{2\pi a^2}$

Sol: $\oint \vec{E}_r \cdot d\vec{A} = \frac{q_{\text{enclosed}}}{\epsilon_0} r$ ----(1)

$$q_{\text{enclosed}}(r) = \int_a^r 4\pi r^2 dr \times \rho$$

$$= \int_a^r 4\pi r^2 dr \times \frac{A}{r}$$

$$= 4\pi A \frac{r^2}{2} \Big|_a^r$$

$$= 2\pi A(r^2 - a^2)$$

$$(1) \Rightarrow E_r \times 4\pi r^2 = \frac{2\pi A(r^2 - a^2)}{\epsilon_0}$$

$$E_r = \frac{A}{2\epsilon_0} \left(1 - \frac{a^2}{r^2}\right)$$

Total E at r = E_r due to volume charge +
E due to charge at centre

$$E_{\text{total}} = \frac{A}{2\epsilon_0} \left(1 - \frac{a^2}{r^2}\right) + \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$

For r² term to vanish $A = \frac{Q}{2\pi a^2}$

13. A combination of capacitors is set up as shown in the figure. The magnitude ----

Ans: 420 N/C

Sol: Charge on 4 μF is 24 μC
Charge on 9 μF is 18 μC
∴ Q = 42 μC

$$E = 9 \times 10^9 \times \frac{42 \times 10^{-6}}{(30)^2} = 420 \text{ N/C}$$

14. The temperature dependence of resistances of Cu and undoped Si in the ----

Ans: Linear increase for Cu, exponential decreases of Si

Sol: Intrinsic carrier concentration

$$n = n_0 e^{-E_g/2k_b T}$$

n increases exponentially with increase in T. Hence conductivity increases exponentially or resistivity decreases exponentially.

For Cu, R = R₀(1 + αt), it is a linear increase.

15. Two identical wires A and B, each of length 'ℓ' carry the same current I. Wire A is bent ----

Ans: $\frac{\pi^2}{8\sqrt{2}}$

Sol: $B_A = \frac{\mu_0 I}{2R}$ (For circle)

$$B_B = \frac{\mu_0 I}{4\pi \frac{a}{2}} \left[\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \right] \times 4 \text{ sides}$$

$$= \frac{\mu_0 I}{\pi a} 2\sqrt{2}$$

[Using $\frac{\mu_0 I}{4\pi d} (\sin\theta_1 + \sin\theta_2)$]

Also ℓ = 2πR = 4a

$$\frac{B_A}{B_B} = \frac{\pi^2}{8\sqrt{2}}$$

16. Hysteresis loops for two magnetic materials A and B ----


Ans: B for electromagnets and transformers

Sol: For transformers to reduce hysteresis loss, area of the loop should be small and for electromagnets the material should have low retentivity and co-ercivity.

17. An arc lamp requires a direct current of 10 A at 80 V to function. If it is ----

Ans: 0.065 H

Sol: I = 10 A
V = 220 V, f = 50 Hz
R = 0.8 Ω
X_L = Lω



$$I = 10 = \frac{220}{\sqrt{64 + L^2 10^5}}$$

$$64 + 10^5 L^2 = 484$$

$$10^5 L^2 = 420$$

$$L^2 = 420 \times 10^{-5}$$

$$= 42 \times 10^{-4}$$

$$L = 6 \times 10^{-2} \\ = 0.06$$

$$N_2 = \frac{3}{4} N_0$$

$$\therefore \text{ratio is } \frac{5}{4}$$

18. Arrange the following electromagnetic radiations per quantum----

Ans: D, B, A, C

Sol: Least : Radiowave
: yellow (500 nm)
: Blue (400 nm)
Highest : X-ray
D → B → A → C

19. An observer looks at a distant tree of height 10 m with a ----

Ans: 20 times nearer

Sol: Theoretical

20. The box of a pin hole camera, of length L, has a hole of radius a. It is assumed that ----

$$\text{Ans: } a = \sqrt{\lambda L} \text{ and } b_{\min} = \sqrt{4\lambda L}$$

Sol: Fresnel distance

$$= \frac{a^2}{\lambda} = L$$

$$a = \sqrt{\lambda L}$$

$$\beta = b_{\min} = \frac{2\lambda L}{\sqrt{\lambda L}} = \sqrt{4\lambda L}$$

21. Radiation of wavelength λ , is incident on a photocell. The fastest emitted----

$$\text{Ans: } v' > v \left(\frac{4}{3}\right)^{1/2}$$

$$\text{Sol: } v = \sqrt{\frac{2}{m} \left(\frac{hc}{\lambda} - \phi_0 \right)}$$

$$v' = \sqrt{\frac{2}{m} \left(\frac{4hc}{3\lambda} - \phi_0 \right)}$$

$$\therefore v' > v \left(\frac{4}{3}\right)^{1/2}$$

22. Half-lives of two radioactive elements A and B are 20 minutes and 40 minutes ----

Ans: 5 : 4

$$\text{Sol: } \frac{N_1}{N_0} = \left(\frac{1}{2}\right)^{\frac{80}{20}} = \left(\frac{1}{2}\right)^4 = \frac{1}{16}$$

$$\frac{N_2}{N_0} = \left(\frac{1}{2}\right)^{\frac{80}{40}} = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$$

$$\Rightarrow \text{decayed } N_1 = \frac{15}{16} N_0$$

23. If a, b, c, d are inputs to a gate and x is its output, then, as per ----

Ans: OR

Sol: Theoretical

24. Choose the correct statement:

(1) In amplitude modulation the ----

Ans: In amplitude modulation the amplitude of the high frequency carrier wave is made to vary in proportion to the amplitude of the audio signal

Sol: Theoretical

25. A screw gauge with a pitch of 0.5 mm and a circular scale with 50 divisions is used to ----

Ans: 0.80 mm

$$\text{Sol: } LC = \frac{1}{100} \text{ mm} \quad ZE = 0.05$$

$$TR = 0.5 + (0.25 + 0.05) \\ = 0.80 \text{ mm}$$

26. A pipe open at both ends has a fundamental frequency f in air. The pipe is dipped ----

Ans: f

$$\text{Sol: } f_o = \frac{v}{2l}$$

$$f_c = \frac{v}{4\left(\frac{l}{2}\right)}$$

$$f_c = f_o = f$$

27. A galvanometer having a coil resistance of 100 Ω gives a full scale deflection, when ----

Ans: 0.01

$$\text{Sol: } S = \frac{I_g G}{I - I_g} \approx \frac{1 \times 10^{-3} \times 100}{10}$$

$$= 0.01 \Omega$$

28. In an experiment for determination of refractive index of glass of prism ----

Ans: 1.5

$$\text{Sol: } i_1 + i_2 = 114$$

$$D = i_1 + i_2 - A \Rightarrow A = 74$$

$$\mu = \frac{\sin \frac{A+D}{2}}{\sin \frac{A}{2}} = \frac{\sin 57}{\sin 37}$$

$$\approx 1.5$$

29. Identify the semiconductor devices whose characteristics are given ----

Ans: Simple diode, zener diode, solar cell, light dependent resistance

Sol: Theoretical

30. For a common emitter configuration, if α and β have their usual meanings, ----

Ans: $\alpha = \frac{\beta}{1-\beta}$

Sol: $I_C = I_C + I_B$

$$\alpha = \frac{I_C}{I_E} \quad \beta = \frac{I_C}{I_B}$$

PART – B – CHEMISTRY

31. At 300 K and 1 atm, 15 mL of a gaseous hydrocarbon requires ----

Ans: C_3H_8

Sol: $C_3H_{8(g)} + 5O_{2(g)} \rightarrow 3CO_{2(g)} + 4H_2O_{(l)}$
 15 mL 75 mL 45 mL
 After combustion, the gases occupy 345 mL

32. Two closed bulbs of equal volume (V) containing an ideal gas initially at pressure ----

Ans: $2P_1 \left(\frac{T_2}{T_1 + T_2} \right)$

Sol: $n_i = n_f$
 $\frac{P_i}{T_1} + \frac{P_i}{T_1} = \frac{P_f}{T_1} + \frac{P_f}{T_2}$
 $\frac{2P_i}{T_1} = \frac{P_f(T_1 + T_2)}{T_1 T_2}$
 $P_f = 2P_i \left(\frac{T_2}{T_1 + T_2} \right)$

33. A stream of electrons from a heated filament was passed between two charged ----

Ans: $\sqrt{2m eV}$

Sol: $\lambda = \frac{h}{\sqrt{2m eV}}$
 $\frac{h}{\lambda} = \sqrt{2m eV}$

34. The species in which the N atoms is in a state of sp hybridisation is: ----

Ans: NO_2^+

Sol: $\begin{matrix} & O \\ & // \\ \oplus N & \\ & \backslash \\ & O \end{matrix}$
 bp = 2
 lp = 0
 Hybridisation = sp

35. The heats of combustion of carbon and carbon monoxide are -393.5 and -283.5 kJ mol⁻¹ ----

Ans: -110.5

Sol: $C_{(s)} + O_{2(g)} \rightarrow CO_{2(g)}$
 $\Delta H = (-393.5) - (-283.5)$
 $= -110$ kJ

36. 18 g of glucose ($C_6H_{12}O_6$) is added to 178.2 g water. The vapour pressure of water (in torr) ----

Ans: 752.4

Sol: $\frac{P^\circ - P_s}{P^\circ} = \frac{n_2}{n_1 + n_2}$
 $\frac{760 - p_s}{760} = \frac{0.1}{0.1 + 9.9}$
 $P_s = 760 - 7.6$
 $= 752.4$ Torr

37. The equilibrium constant at 298 K for a reaction $A + B \rightleftharpoons C + D$ is 100. ----

Ans: 1.818

Sol: $A + B \rightleftharpoons C + D$

Initial	1	1	1	1
Eqbm	1-x	1-x	1+x	1+x

 $K_c = \frac{(1+x)^2}{(1-x)^2}$
 $10 = \frac{1+x}{1-x}$
 $x = \frac{9}{11}$
 $[D] = 1+x = 1 + \frac{9}{11}$
 $= \frac{20}{11}$
 $= 1.818$


38. Galvanization is applying a coating of: ----

Ans: Zn

Sol: Coating of Zn on metal surface is galvanization.

39. Decomposition of H_2O_2 follows a first order reaction. ----

Ans: 6.93×10^{-4} mol min⁻¹



Rate of formation of oxygen
 $= \frac{0.693}{25} \times 0.05 \times \frac{1}{2}$
 $= 6.93 \times 10^{-4} \text{ mol min}^{-1}$

40. For a linear plot of $\log(x/m)$ versus $\log p$ in a Freundlich adsorption isotherm, ----

Ans: Only $1/n$ appears as the slope

Sol: $\frac{x}{m} = k P^{1/n}$
 $\log \frac{x}{m} = \log k + \frac{1}{n} \log P$
 $\frac{1}{n}$ is the slope

41. Which of the following atoms has the highest first ionization energy? ----

Ans: Sc

Sol: Sc has the highest first ionisation enthalpy
Sc – 631 kJ mol^{-1}
Na – 496 kJ mol^{-1}
K – 419 kJ mol^{-1}
Rb – 403 kJ mol^{-1}

42. Which one of the following ores best concentrated by froth floatation method? ----

Ans: Galena

Sol: Galena is PbS .
Sulphide ores are concentrated by froth floatation method.

43. Which one of the following statements about water is **FALSE**? ----

Ans: There is extensive intramolecular hydrogen bonding in the condensed phase

Sol: There is no intramolecular hydrogen bonding in water.


44. The main oxides formed on combustion of Li, Na and K in excess of air are, respectively----

Ans: Li_2O , Na_2O_2 and KO_2

Sol: Li mainly forms monoxide, Li_2O , Na forms peroxide, Na_2O_2 and K forms superoxide, KO_2 .

45. The reaction of Zinc with dilute and concentrated nitric acid, respectively ----

Ans: N_2O and NO_2



46. The pair in which phosphorus atoms have a formal oxidation state of +3 is:----

Ans: Orthophosphorus and pyrophosphorus acids

Sol: Orthophosphorus acid is H_3PO_3 and pyrophosphorus acid is $\text{H}_4\text{P}_2\text{O}_5$.

47. Which of the following compounds is metallic and ferromagnetic? ----

Ans: CrO_2

Sol: CrO_2 is metallic and ferromagnetic

48. The pair having the same magnetic moment is: ----

Ans: $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$ and $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$

Sol: $\text{Cr}^{2+} \rightarrow [\text{Ar}]3d^4$
 $\text{Fe}^{2+} \rightarrow [\text{Ar}]3d^6$
Both contains 4 unpaired electrons.

49. Which one of the following complexes shows optical isomerism? ----

Ans: $\text{cis}[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$

Sol: $\text{cis}[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$ is optically active.

50. The concentration of fluorides, lead, nitrate and iron in a water sample ----

Ans: Nitrate



Sol: The maximum permissible concentration of NO_3^- in drinking water is only 50 ppm.

51. The distillation technique most suited for separating glycerol from spent-lye----

Ans: Distillation under reduced pressure

Sol: Glycerol is purified by distillation under reduced pressure.

52. The product of the reaction given below is:----



53. The absolute configuration of ----

Ans: (2S, 3R)

Sol: According to CIP rules.


54. 2-chloro-2-methylpentane on reaction with sodium methoxide in methanol yields: ----

Ans: (c) only

Sol: Tertiary alkyl halide will undergo elimination with NaOCH₃. No ether will be formed.


55. The reaction of propene with HOCl (Cl₂ + H₂O) proceeds through the intermediate ----

Ans: CH₃-CH⁺-CH₂-Cl



56. In the Hoffmann bromamide degradation reaction, the number of moles of ----

Ans: Four moles of NaOH and one mole of Br₂




57. Which of the following statements about low density polythene is FALSE?----

Ans: It is used in the manufacture of buckets, dust-bins etc

Sol: HDP is used for the manufacture of buckets, dust-bins, etc

58. Thiol group is present in ----

Ans: Cysteine



59. Which of the following is the anionic detergent? ----

Ans: Sodium lauryl sulphate

Sol: Sodium lauryl sulphate is an anionic detergent

60. The hottest region of Bunsen flame ----

Ans: region 2

Sol: Region 2 forms the hottest part of Bunsen flame.

PART - C - MATHEMATICS

61. If $f(x) + 2f\left(\frac{1}{x}\right) = 3x$, $x \neq 0$, and $S = \{x \in \mathbf{R} : f(x) \neq 0\}$ ----

Ans: contains exactly two elements

Sol: Let $f(x) = A$ and $f\left(\frac{1}{x}\right) = B$

Given : $A + 2B = 3x$, replacing x by $\frac{1}{x}$

$$\Rightarrow 2B + 4A = \frac{6}{x}$$

$$\therefore 3A = \frac{6}{x} - 3x$$

$$\Rightarrow A = \frac{2}{x} - x$$

$$\frac{2}{x} - x = \frac{-2}{x} + x$$

$$2x = \frac{4}{x}$$

$$x^2 = 2$$

$$\therefore x = \pm\sqrt{2} \Rightarrow \text{Two elements}$$

62. A value of θ for which $\frac{2 + 3i \sin \theta}{1 - 2i \sin \theta}$ ----

Ans: $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$

Sol: $\frac{2 + 3i \sin \theta}{1 - 2i \sin \theta}$ purely imaginary

$$\Rightarrow \frac{2 - 6 \sin^2 \theta}{1 + 4 \sin^2 \theta} = 0$$

$$\Rightarrow \sin^2 \theta = \frac{1}{3}$$

$$\Rightarrow \sin \theta = \frac{1}{\sqrt{3}}$$

$$\theta = \sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

63. The sum of all real values of x satisfying the equation----

Ans: 3

Sol: $x^2 - 5x + 5 = 1$ and $x^2 + 4x - 60 \in \mathbb{R}$
 Or $x^2 - 5x + 5 \in \mathbb{R}$ and non zero and $x^2 + 4x - 60 = 0$
 or $x^2 - 5x + 5 = -1$ and $x^2 + 4x - 60$ is even
 $\therefore x = 1, 4$ or $x = -10, 6$ or $x = 2, 3$
 But $x = 3 \Rightarrow x^2 + 4x - 60$ is odd
 $\therefore x = 1, 4, -10, 6, 2$
 Sum of the roots = 3

64. If $A = \begin{bmatrix} 5a & -b \\ 3 & 2 \end{bmatrix}$ and $A \text{ adj } A = AA^T$, then----

Ans: 5

Sol: $A \text{ adj } A = AA^T$
 $|A|I = AA^T$
 $\Rightarrow \begin{bmatrix} 10a+3b & 0 \\ 0 & 10a+3b \end{bmatrix} = \begin{bmatrix} 25a^2-b^2 & 15a-2b \\ 15a-2b & 13 \end{bmatrix}$
 $\Rightarrow 10a+3b = 13$
 $15a-2b = 0$
 Solving $a = \frac{2}{5}$ $b = 3$
 $\therefore 5a + b = 2 + 3 = 5$

65. The system of linear equations

$$\begin{aligned} x + \lambda y - z &= 0 \\ \lambda x - y - z &= 0 \\ x + y - \lambda z &= 0 \end{aligned}$$

Ans: exactly three values of λ

Sol: $\begin{vmatrix} 1 & \lambda & -1 \\ \lambda & -1 & -1 \\ 1 & 1 & -\lambda \end{vmatrix} = 0$
 $1(\lambda + 1) - \lambda(-\lambda^2 + 1) - 1(\lambda + 1) = 0$
 $\lambda^3 - \lambda = 0 \Rightarrow \lambda(\lambda^2 - 1) = 0$
 $\lambda = 0, \lambda^2 = 1 \Rightarrow \lambda = \pm 1$
 Exactly three values of λ

66. If all the words (with or without meaning) having five letters----

Ans: 58^{th}

Sol: Arranging SMALL is ascending order ALLMS

$$\text{Starting with A} - \frac{4!}{2!} = \frac{24}{2} = 12$$

$$\text{Starting with L} - 4! = 24$$

$$\text{Starting with M} - \frac{4!}{2!} = \frac{24}{2} = 12$$

$$\text{Starting with SA} - \frac{3!}{2!} = \frac{6}{2} = 3$$

$$\text{Starting with SL} - 3! = 6$$

$$\text{SMALL} \Rightarrow 1$$

$$\text{Total} = 12 + 24 + 12 + 3 + 6 + 1 = 58^{\text{th}}$$

67. If the number of terms in the expansion of ----

Ans: 729

Sol: Number of terms = 28

$$\frac{(n+1)(n+2)}{2} = 28$$

$$\Rightarrow n + 1 = 7 \Rightarrow n = 6$$

Put $x = 1$

$$(1 - 2 + 4)^6 = 3^6 = 729$$

68. If the 2^{nd} , 5^{th} , and 9^{th} terms of a non - constant A.P are in G.P----

Ans: $\frac{4}{3}$

Sol: $a + d, a + 4d, a + 8d$ are in GP.

$$\therefore (a + 4d)^2 = (a + d)(a + 8d)$$

$$a^2 + 16d + 8ad = a^2 + 8d^2 + 9ad$$

$$\Rightarrow 8d^2 - ad = 0$$

$$d(8d - a) = 0 \quad d \neq 0$$

$$\therefore a = 8d$$

\therefore Terms are $9d, 12d, 16d$ are in G.P

$$\therefore r = \frac{12d}{9d} = \frac{4}{3}$$

69. If the sum of the first ten terms of the series ----

Ans: 101

Sol: The sequence may be rearranged as

$$\left(\frac{8}{5}\right)^2 + \left(\frac{12}{5}\right)^2 + \left(\frac{16}{5}\right)^2 \dots \text{up to ten terms}$$

$$= \frac{16}{25} [2^2 + 3^2 + 4^2 + \dots + 11^2]$$

$$= \frac{16}{25} \left[\frac{11 \times 12 \times 23}{6} - 1 \right] = \frac{16}{25} \times 505$$

$$= \frac{16}{5} m(\text{given}) \Rightarrow m = 101$$

70. Let $p = \lim_{x \rightarrow 0^+} (1 + \tan^2 \sqrt{x})^{\frac{1}{2x}}$ then $\log p$ ----

Ans: $\frac{1}{2}$

$$\text{Sol: } p = e^{\lim_{x \rightarrow 0} \left(\frac{\tan^2 \sqrt{x}}{2x} \right)}$$

$$= e^{\lim_{x \rightarrow 0} \frac{2 \tan \sqrt{x} \sec^2 \sqrt{x} \cdot \frac{1}{2\sqrt{x}}}{2}}$$

$$= e^{\lim_{x \rightarrow 0} \left(\frac{\tan \sqrt{x}}{\sqrt{x}} \right) \sec^2 \sqrt{x}} \frac{1}{2}$$

$$= e^{\frac{1}{2}}$$

$$\therefore \log p = \frac{1}{2}$$

71. For $x \in \mathbf{R}$, $f(x) = |\log 2 - \sin x|$ and $g(x) = \dots$

Ans: $g'(0) = \cos(\log 2)$

Sol: $g(x) = f(f(x))$
 $g'(0) = f'(f(0)) \cdot f'(0)$
near $x = 0$, $f(x) = |\log 2 - \sin x| = \log 2 - \sin x$
 $f'(0) = -\cos 0 = -1$
 $f(0) = |\log 2| = \log 2$
 $\therefore f'(f(0)) = f'(\log 2) = -\cos(\log 2)$
 $\therefore g'(0) = [-\cos(\log 2)] \cdot (-1) = \cos(\log 2)$

72. Consider $f(x) = \tan^{-1} \dots$

Ans: $\left(0, \frac{2\pi}{3} \right)$

Sol: $f(x) = \tan^{-1} \frac{1 + \cos\left(\frac{\pi}{2} - x\right)}{1 - \cos\left(\frac{\pi}{2} - x\right)}$

$$= \tan^{-1} \cot\left(\frac{\pi}{4} - \frac{x}{2}\right)$$

$$= \frac{\pi}{4} + \frac{x}{2}$$

$$f'(x) = \frac{1}{2}$$

\therefore Slope of normal = -2

\therefore Equation $y - \frac{\pi}{3} = -2\left(x - \frac{\pi}{6}\right)$

Satisfying $\left(0, \frac{2\pi}{3} \right)$

73. A wire of length 2 units is cut into two parts which are bent respectively----

Ans: $x = 2r$

Sol: Given $4x + 2\pi r = 2$

$$\Rightarrow x = \frac{1 - \pi r}{2}$$

$$\text{Area} = x^2 + \pi r^2$$

$$= \frac{1 + \pi^2 r^2 - 2\pi r}{4} + \pi r^2$$

$$\frac{dA}{dr} = 0 \Rightarrow r = \frac{1}{\pi + 4}$$

$$x = \frac{2}{\pi + 4}$$

$$\frac{d^2A}{dr^2} > 0$$

$$\therefore x = 2r$$

74. The integral $\int \frac{2x^{12} + 5x^9}{(x^5 + x^3 + 1)^3} dx$ is equal to----

Ans: $= \frac{x^{10}}{2(x^5 + x^3 + 1)^2} + C$

Sol: $\int \frac{(2x^{12} + 5x^9) dx}{(x^5 + x^3 + 1)^3} = \int \frac{\left(\frac{2}{x^3} + \frac{5}{x^6}\right) dx}{\left(1 + \frac{1}{x^2} + \frac{1}{x^5}\right)^3}$

$$= \int \frac{-du}{u^3} \text{ if } u = 1 + \frac{1}{x^2} + \frac{1}{x^5}$$

$$= \frac{1}{2u^2} + C$$

$$= \frac{1}{2\left(1 + \frac{1}{x^2} + \frac{1}{x^5}\right)^2} + C$$

$$= \frac{x^{10}}{2(x^5 + x^3 + 1)^2} + C$$

75. $\lim_{n \rightarrow \infty} \left(\frac{(n+1)(n+2)\dots 3n}{n^{2n}} \right)^{\frac{1}{n}}$ is equal to----

Ans: $\frac{27}{e^2}$

Sol: $L = \lim_{n \rightarrow \infty} \left[\frac{(n+1)(n+2)\dots(n+2n)}{n^{2n}} \right]^{\frac{1}{n}}$

$$\log L = \lim_{n \rightarrow \infty} \frac{1}{n} \left[\log\left(1 + \frac{1}{n}\right) + \log\left(1 + \frac{2}{n}\right) + \dots \right. \\ \left. \dots + \log\left(1 + \frac{2n}{n}\right) \right]$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^{2n} \log\left(1 + \frac{r}{n}\right)$$

$$= \int_0^2 \log(1+x) dx$$

$$= \left\{ \log(1+x) \times x \Big|_0^2 - \int_0^2 \frac{1}{1+x} dx \right\}$$

$$= 2\log 3 - (2 - \log 3)$$

$$= 3\log 3 - 2$$


$$= \log\left(\frac{27}{e^2}\right)$$

$$L = \frac{27}{e^2}$$

76. The area (in sq. units) of the region ----

Ans: $\pi - \frac{8}{3}$

Sol:



$$y^2 = 2x \quad (1)$$

$$x^2 - y^2 - 4x = 0 \quad (2)$$

To find the x- coordinates of their points of intersection, solve (1) and (2)

$$x^2 + 2x - 4x = 0$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x = 0 \text{ or } 2$$

$$\therefore \text{Area} = \int_0^2 (y_1 - y_2) dx$$

$$= \int_0^2 [\sqrt{4 - (x-2)^2} - \sqrt{2} \sqrt{x}] dx$$

$$= \left[\frac{x-2}{2} \sqrt{4 - (x-2)^2} + \frac{4}{2} \sin^{-1} \left(\frac{x-2}{2} \right) - \sqrt{2} \frac{2}{3} x^{3/2} \right]_0^2$$

$$= \pi - \frac{8}{3}$$

77. If a curve $y = f(x)$ passes through the point $(1, -1)$ and satisfies----

Ans: $\frac{4}{5}$

Sol: $y(1 + xy)dx = xdy$

$$x \frac{dy}{dx} = y + xy^2$$

$$-\frac{1}{y^2} \frac{dy}{dx} = \frac{-1}{xy} - 1$$

$$-\frac{1}{y^2} \frac{dy}{dx} + \frac{1}{xy} = -1$$

i.e. $\frac{dY}{dx} + \frac{Y}{x} = -1$ where $Y = \frac{1}{y}$

Solution is $Yx = \frac{-x^2}{2} + C$

$$-1 = -\frac{1}{2} + C \Rightarrow C = -\frac{1}{2}$$

$$\frac{x}{y} = -\frac{x^2 + 1}{2}$$

$$y = \frac{-2x}{1+x^2}; y\left(-\frac{1}{2}\right) = \frac{1}{1+\frac{1}{4}}$$

$$= \frac{4}{5}$$

78. Two sides of a rhombus are along the lines, $x - y + 1 = 0$ ----

Ans: $\left(\frac{1}{3}, \frac{-8}{3}\right)$

Sol: Point of intersection of $x - y + 1 = 0$ and $7x - y - 5 = 0$ is $(1, 2)$ is one of the vertex. The image point is also not the required point. Equation of the diagonal containing $(1, 2)$ and $(-1, -2)$ is $2x - y + 1 = 0$

\therefore The other diagonal is $x + 2y + k = 0$
i.e. $-1 - 4 + k = 0$
 $x + 2y + 5 = 0$

Solving $x + 2y + 5 = 0$ and $7x - y - 5 = 0$

We get $x = \frac{1}{3}$ and $x = \frac{-8}{3}$

\therefore required vertex is $\left(\frac{1}{3}, \frac{-8}{3}\right)$

79. The centres of those circles which touch the circle, $x^2 + y^2 - 8x - 8y - 4 = 0$ ----

Ans: a parabola

Sol: If (a, b) is the centre then radius is b (if $b > 0$) & $-b$ (if $b < 0$)


We have
 $(a - 4)^2 + (b - 4)^2 = (6 + b)^2$ if $b > 0$
 $(a - 4)^2 + (b - 4)^2 = (6 - b)^2$ if $b < 0$

\therefore locus is
 $(x - 4)^2 + (y - 4)^2 = (6 + y)^2$ or
 $(x - 4)^2 + (y - 4)^2 = (6 - y)^2$, each of which is a parabola

80. If one of the diameters of the circle, given by the equation, $x^2 + y^2 - 4x + 6y - 12 = 0$ ----

Ans: $5\sqrt{3}$

Sol:



AB is the diameter of the circle $x^2 + y^2 - 4x + 6y - 12 = 0$ ---- (1)

Midpoint of AB = centre of (1) = $(2, -3)$

CB = radius of (1)
 $= \sqrt{4 + 9 + 12} = 5$

$OC = \sqrt{5^2 + 5^2} = 5\sqrt{2}$

\therefore Radius of S = OB
 $= \sqrt{OC^2 + CB^2}$
 $= \sqrt{25 + 50}$

$$= \sqrt{75}$$

$$= 5\sqrt{3}$$

81. Let P be the point on the parabola, $y^2 = 8x$ which is at a minimum distance from the centre C of the circle----

Ans: $x^2 + y^2 - 4x + 8y + 12 = 0$

Sol: Let P be the point $(2t^2, 4t)$
 C is $(0, -6)$
 $D = PC^2 = (2t^2)^2 + (4t + 6)^2$
 $= 4t^4 + 16t^2 + 48t + 36$

$$\frac{d}{dt}(D) = 16t^3 + 32t + 48$$

$$\frac{d^2}{dt^2}(D) = 48t^2 + 32$$

$$\frac{d}{dt}(D) = 0 \Rightarrow t = -1$$

\therefore P is $(2, -4)$

\therefore Circle with centre at P is
 $(x - 2)^2 + (y + 4)^2 = r^2$
 Since, passes through $(0, -6)$
 $r^2 = 8$

\therefore Required circle is
 $(x - 2)^2 + (y + 4)^2 = 8$
 or $x^2 + y^2 - 4x + 8y + 12 = 0$

82. The eccentricity of the hyperbola whose length of the latus rectum is equal to 8 ----

Ans: $\frac{2}{\sqrt{3}}$

Sol: $\frac{2b^2}{a} = 8$ ---- (1)

$2b = ae$ ---- (2)

(2) $\Rightarrow 4b^2 = a^2e^2$

$4a^2(e^2 - 1) = a^2e^2$

$\Rightarrow 4e^2 - 4 = e^2$

$\Rightarrow e^2 = \frac{4}{3}$

$\Rightarrow e = \frac{2}{\sqrt{3}}$

83. The distance of the point $(1, -5, 9)$ from the plane $x - y + z = 5$ measured along the ----

Ans: $10\sqrt{3}$

Sol: $(1, -5, 9)$
 $x - y + z = 5$
 $x = y = z$

$$\frac{x-1}{1} = \frac{y+5}{1} = \frac{z-9}{1} = \lambda \Rightarrow$$

Point is $(\lambda + 1, \lambda - 5, \lambda + 9)$

Where the two meet the plane

$\lambda + 1 - \lambda + 5 + \lambda + 9 = 5$

$\lambda = -10$

\therefore point is $(-9, -15, 1)$

distance = $\sqrt{100 + 100 + 100} = 10\sqrt{3}$

84. If the line $\frac{x-3}{2} = \frac{y+2}{-1} = \frac{z+4}{3}$ lies in----

Ans: 2

Sol: $\frac{x-3}{2} = \frac{y+2}{-1} = \frac{z+4}{3}$

lies on $\ell x + m y - z = 9$

$3\ell - 2m + 4 = 9$

$2\ell - m - 3 = 0$

$\Rightarrow 3\ell - 2m = 5$

$2\ell - m = 3$

$\Rightarrow \ell = 1, m = -1$

$\ell^2 + m^2 = 2$

85. Let \bar{a} , \bar{b} and \bar{c} be three ----

Ans: $\frac{5\pi}{6}$

Sol: $\frac{\sqrt{3}}{2}(\bar{b} + \bar{c}) = \bar{a} \times (\bar{b} \times \bar{c})$

$= (\bar{a} \cdot \bar{c})\bar{b} - (\bar{a} \cdot \bar{b})\bar{c}$

$\therefore (\bar{a} \cdot \bar{b}) = -\frac{\sqrt{3}}{2}$

\therefore angle between \bar{a} & \bar{b} is $\frac{5\pi}{6}$

(\bar{a} , \bar{b} are unit vectors)

86. If the standard deviation of the numbers 2, 3, a and 11 is 3.5, ----

Ans: $3a^2 - 32a + 84 = 0$

Sol: Variance $\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2$

$12.25 = \frac{4+9+121+a^2}{4} - \frac{(16+a)^2}{16}$

$= \frac{134 \times 4 + 4a^2 - 256 - 32a - a^2}{16}$

$\Rightarrow 196 = 3a^2 - 32a + 536 - 256$

$\Rightarrow 3a^2 - 32a + 84 = 0$

87. Let two fair six - faced dice A and B be thrown simultaneously, ----

Ans: E_1, E_2, E_3 are independent

Sol: $E_1 = \{(4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6)\}$

$E_2 = \{(1, 2), (2, 2), (3, 2), (4, 2), (5, 2), (6, 2)\}$

$E_3 = \{(1, 2), (1, 4), (1, 6), (2, 1), (2, 3), (2, 5), (3, 2), (3, 4), (3, 6), (4, 1), (4, 3), (4, 5), (5, 2), (5, 4), (5, 6), (6, 1), (6, 3), (6, 5)\}$

$$P(E_1) \cdot P(E_2) = \frac{1}{6} \cdot \frac{1}{6} = \frac{1}{36} = P(E_1 \cap E_2)$$

$\therefore E_1, E_2$ independent

$$P(E_1) P(E_3) = \frac{1}{6} \cdot \frac{1}{2} = \frac{1}{12} = P(E_1 \cap E_3)$$

E_1, E_3 independent

$$P(E_2) P(E_3) = \frac{1}{6} \cdot \frac{1}{2} = \frac{1}{12} = P(E_2 \cap E_3)$$

$$P(E_1) P(E_2) P(E_3) = \frac{1}{6} \cdot \frac{1}{6} \cdot \frac{1}{2} = \frac{1}{72} \neq P(E_1 \cap E_2 \cap E_3)$$

$\therefore E_1, E_2, E_3$ not independent

88. If $0 \leq x < 2\pi$, then the number of real values of x , which satisfy the equation----

Ans: 7

Sol: The equation $\cos x (\cos 2x + \cos 3x) = 0$

$$\therefore \cos x = 0 \text{ or } \cos 3x = -\cos 2x$$

$$x = \frac{\pi}{2} \text{ or } \frac{3\pi}{2} \text{ or } 3x = (2n+1)\pi \pm 2x$$

$$\Rightarrow x = (2n+1)\pi$$

$$\text{or } x = (2n+1)\frac{\pi}{5}$$


$$\text{Solution are } \frac{\pi}{5}, \frac{\pi}{2}, \frac{3\pi}{5}, \pi, \frac{7\pi}{5}, \frac{9\pi}{5}, \frac{3\pi}{2}$$

Total 7 nos

89. A man is walking towards a vertical pillar in a straight path, at a uniform speed----

Ans: 5

Sol:



OT is tower – height h (say)

AB = x

OB = y

$$x + y = \frac{h}{\tan 30^\circ} = h\sqrt{3}$$

$$x = \frac{h}{\tan 60^\circ} = \frac{h}{\sqrt{3}}$$

$$\therefore y = \frac{2h}{\sqrt{3}}$$


$$\frac{y}{x} = \frac{1}{2}$$

x takes 10 m, \therefore true for $y = 5$ m

90. The Boolean Expression $(p \wedge \sim q) \vee q \vee (\sim p \wedge q)$

Ans: $p \vee q$

Sol:



$$(p \wedge \sim q) \vee q \vee (\sim p \wedge q) = p \vee q$$